

Kindergarten through 12th Grade Immunization Chart

Immunizations REQUIRED

VACCINE	Number of Doses	Grades K-12 (4 – 18 years of age)
		Vaccines must follow MINIMUM INTERVALS and AGES to be valid. A 4 day grace period applies in most situations.
Rotavirus (2 dose series)	2-6 mos	RV1 2 dose schedule RV5 3 dose schedule
Diphtheria/Tetanus/ Pertussis (DTaP or DT) <i>Only licensed through 6 years of age</i>	4 to 5	5 DTaP or DT unless dose 4 given on or after the 4 th b-day. Final dose of DTaP or DT must be given no sooner than 4 years of age
Tetanus/Diphtheria/ Pertussis <i>For students 7 years of age or older who did not have full series of DTaP or DT</i>	3 or 4	3 doses Tetanus/diphtheria containing vaccines (DTaP, DT, Td, Tdap) is required. 4 doses required if 1 st dose of DTaP or DT is given before 1 year of age. 1 dose of Tdap is to be given if DTaP series not completed and student is at least 7 years of age. Tdap is required at 6 th grade entry thru 12 th grade.
Haemophilus influenza type b (HIB)	1-4 doses depending on age	Dosage and catch up schedule is complex. Please refer to CDC catch up schedule.
Pneumooccal	2-4 doses	Dosage and catch up schedule is complex. Please refer to CDC catch up schedule
Polio (IPV) <i>With combination of OPV & IPV, need series of 4 doses</i>	3 to 4	4 IPV unless 3 rd dose is given on or after 4 th birthday. Students who were compliant with 3 or 4 doses. (4 weeks minimum intervals between doses) prior to August 7, 2009 have met the requirement.
Measles/Mumps/Rubella (MMR) <i>There must be at least a 28 day interval between 2 live vaccines.</i>	2	The 1 dose is not valid if administered more than 4 days before the 1 st birthday. 2 doses are required for students entering Kindergarten and through 12 th grade.
Varicella (Chickenpox) <i>There must be at least a 28 day interval between 2 live vaccines.</i>	2	The 1 st dose is not valid if administered more than 4 days before the 1 st birthday. 2 doses are required for students entering Kindergarten and through 12 th grade. <i>Note: no vaccine required if there is laboratory documentation of chickenpox disease or a disease screening by a healthcare provider has been performed.</i>
Hepatitis B <i>Dosing must follow intervals between doses and last dose must be administered at or over 24 wks of age.</i>	3	The 2 nd dose must be administered at least 4 weeks after the first dose. The 3 rd dose must be administered at least 16 weeks after the 1 st dose, at least 8 weeks after the 2 nd dose, and the final dose must be administered no sooner than 24 weeks of age. <i>Note: there is a specific 2-dose series for ages 11-15 years only using only adult vaccine.</i>
Hepatitis A (Hep A)	2	All children 1 year of age and older
Meningococcal Meningitis (MENAcwy)	2	Adolescents 11-18 (11-12, 16-18)

Recommended Vaccines for the best protection against vaccine preventable disease

VACCINE	Number of Doses	Grades K-12 (4 – 18 years of age)
		Vaccines administered < 4 days the minimum age are valid.
Influenza (flu)	1 to 2	2 doses initially if under 9 years of age with a minimum interval of 28 days between doses, then 1 dose annually thereafter. Recommended for all children 6 months of age and older
<u>Meningococcal Meningitis</u> MenB	2 doses	Adolescents 16-18 years
Human Papillomavirus (9vHPV)	2 to 3	Adolescents 11-18 years of age Series initiation age 9-14 2 doses 6-12 mos apart Series initiation 15+ 3 doses 0, 1-2, and 6 mos